

Crown & Bridge Post-Operative Instructions

- While you are numb be careful not to bite your cheek, tongue or your lip. You may want to avoid chewing on this area especially hot foods and liquids until the numbing completely wears off.
- Some irritation of the gum tissue along with the anesthetic injection site may result in some soreness for a few days.

Temporary Crown or Bridge

- You have just received a temporary that will serve you for a short period of time while your custom permanent crown or bridge is being made. This will protect the tooth and hold the tooth in position so the permanent crown or bridge will fit properly.
- The size, shape, and color of the temporary may differ from the permanent crown or bridge.
- Your temporary is not strong. It may break or come off occasionally. Avoid chewing hard or sticky food on your temporary.
- Temporary cement is used so that the temporary can be easily removed at your next appointment. If your temporary comes off between appointments, call us to have it recemented.
- Brush softly and floss carefully around your temporary. When flossing around your temporary, slide the floss out between your teeth rather than snapping up between the teeth. It is important that the gum tissue remains healthy or we may not be able to seat your new crown or bridge.

Permanent Crown or Bridge

- Sensitivity to hot and cold is common for a few weeks after cementing your crown or bridge. If sensitivity does occur, we recommend avoiding extreme temperature change on your new crown or bridge.
- Your crown or bridge may have a different texture than your original tooth and may be contoured slightly different. After a few days your mouth will become completely accustomed to it.
- As with the temporary, if the bite does not feel comfortable, please contact us for a simple adjustment.
- Proper brushing, flossing, and routine dental cleaning & exams are necessary to maintain your final crown or bridge.